

# Phytosociological Studies of Some Medicinal Trees from Dongargan Sacred Grove from Ahmednagar District, Maharashtra, India

Thete AM<sup>1</sup> and Sharma PP<sup>2\*</sup>

<sup>1</sup>Shri.dhokeshwar College, Takalidhokeshwar, Tal- Parner, Dist- Ahmednagar (M.S)

<sup>2</sup>Indraraj Arts, Commerce and Science College, Sillod, Dist- Aurangabad, (M.S)

Email: [anjalkhilari@gmail.com](mailto:anjalkhilari@gmail.com)<sup>1</sup> | [drppsharma6848@gmail.com](mailto:drppsharma6848@gmail.com)<sup>2</sup>

## Manuscript Details

Available online on <https://www.irjse.in>

ISSN: 2322-0015

Editor: Dr. Arvind Chavhan

## Cite this article as:

Thete AM and Sharma PP. Phytosociological Studies of Some Medicinal Trees from Dongargan Sacred Grove from Ahmednagar District, Maharashtra, India, *Int. Res. Journal of Science & Engineering*, 2020, Special Issue A9: 223-226.

Article published in Special issue of International e-Conference on "Emerging trends and Challenges In life sciences" organized by Department of Botany, Indraraj Arts, Commerce & Science College, Sillod-431112, Dist Aurangabad, Maharashtra, India date, June 18-19, 2020.



Open Access This article is licensed under a Creative Commons Attribution 4.0 International License, which permits use, sharing, adaptation, distribution and reproduction in any medium or format, as long as you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons license, and indicate if changes were made. The images or other third party material in this article are included in the article's Creative Commons license, unless indicated otherwise in a credit line to the material. If material is not included in the article's Creative Commons license and your intended use is not permitted by statutory regulation or exceeds the permitted use, you will need to obtain permission directly from the copyright holder. To view a copy of this license, visit <http://creativecommons.org/licenses/by/4.0/>

## Abstract

During the Phytosociological study of Dongargan sacred grove of Ahmednagar district of Maharashtra 20 tree species used for medicinal purpose by the local people are considered. Local people utilize these plants for treating various diseases/ailments. For the purpose of study Quadrante method is used. Quadrates of 100m X 100m size were randomly laid to study the trees species. Analytical characters such as frequency, density and abundance are calculated with the help of standard formulae.

**Keywords:** Colonization, mobile phone, health hazard, diarrhea and dysentery. Sacred grove, Phytosociological Study, Medicinal Plants, Quadrates, Analytical characters

## Introduction

A Sacred grove is a terrestrial segment containing dense flora having local biodiversity, protected by local people or ethnic people in admiration of the deity exist in that grove, this grove is locally called as Deorai. Sacred groves are not only substantial sites for regional biodiversity but also deliver fundamental natural resources to local people. Some local people having ethnic knowledge about the vegetation, which are used as remedy in treating different diseases were referred. The information about dispersal, habitat, demographic features and value of these plant species are

transmitted from one generation [1]. Sacred groves preserve useful plant species which are threatened and endemic due to over exploitation and rapid urbanization. For maintaining the biodiversity, ecological study of sacred grove is necessary. The statistical data gives focus on species, which are dominant in communities to know their dominance certain analytical characters such as Frequency, Density and Abundance of species in a community are expressed in quantity by using quadrature method.

## Material and Method

The current study was carried out in the Dongargan sacred grove for knowing the ecological status of plant species. Quadrature method is used for determination of data. The total area of the sacred grove is 37 acres. Apart

from this sacred forest region dense area of forest used for study. This area was divided into 5 quadrates randomly, 100m x 100m size of single quadrature taken for tree species. The data was collected during rainy season. No. of individual species in each quadrature are recorded and percentage frequency, density and dominance were determined as per Curtis and McIntosh, [2]. The plants are identified with the help of Flora of Ahmednagar district [3]. Counted the number of individuals of each species occur in the quadrature. The frequency of individual species calculated. Abundance of any species is expressed as a percentage of the total number of species present in community and therefore it is a relative measure. In sampling the abundance of species, individual species are counted instead of just noting their presence or absence was done while the frequency of a species.

### Formulas used for calculating Frequency, Density and Abundance:

$$\text{Frequency (\%)} = \frac{\text{Number of quadrates in which the species occurred}}{\text{Total number of quadrates studied.}} \times 100$$

$$\text{Density} = \frac{\text{Total number of individuals of a species in all quadrates}}{\text{Total number of quadrates studied.}}$$

$$\text{Abundance} = \frac{\text{Total number of individuals of a species in all quadrates}}{\text{Total number of quadrates in which the species occurred}}$$

## Results

Table 1: Observations.


Sr. no.	Botanical name of plant species	Individual in a quadrature					Total no of individuals in a quadrature	Total no of quadrates which species Occurred	Density/Unit area	Abundance	Frequency%	Frequency Class
1	<i>Albizia lebeck</i> (L.) Bth.	-	-	-	-	1	01	01	0.2	1.0	20	A
2	<i>Acacia chundra</i> (Roxb.ex Rottle.) Willd.	-	2	-	-	1	03	02	0.6	1.5	40	B
3	<i>Acacia nilotica</i> (L) Willd.	4	7	3	5	3	22	05	4.4	4.4	100	E
4	<i>Aegle marmelos</i> (L.) Corr	2	-	-	1	2	05	03	1.0	1.6	60	C
5	<i>Azadirachta indica</i> A.Juss.	3	2	-	2	3	10	04	2.0	2.5	80	D
6	<i>Bauhinia variegata</i> L.	3	-	4	2	-	09	03	1.8	3.0	60	C

7	<i>Cassia auriculata</i> L.	5	3	3	4	-	15	04	3.0	3.7	80	D
8	<i>Catunaregam spinosa</i> (Thunb) Tirveng	1	-	-	1	-	02	02	0.4	1.0	40	B
9	<i>Citrus limon</i> (L)	-	-	1	-	-	01	01	0.2	1.0	20	A
10	<i>Gmelina arborea</i> Roxb.	2	-	-	-	1	03	02	0.6	1.5	40	B
11	<i>Helicteres isora</i> L.	2	-	-	1	-	03	02	0.6	1.5	40	B
12	<i>Lawsonia inermis</i> L.	-	1	-	-	-	01	01	0.2	1.0	20	A
13	<i>Melia azadirach</i> L.	2	-	-	3	1	06	03	1.2	2.0	60	C
14	<i>Pongamia pinnata</i> (L) Pierre	-	2	3	-	4	09	03	1.8	3.0	60	C
15	<i>Syzygium cumini</i> (L) Skeels	-	-	2	1	1	04	03	0.8	1.3	60	C
16	<i>Tamarindus indica</i> L.	3	2	-	2	3	10	04	2.0	2.5	80	D
17	<i>Tephrosia purpurea</i> (L.)Pers.	2	3	-	-	1	06	03	1.2	2.0	60	C
18	<i>Tectona grandis</i> L. f.	-	2	-	1	-	03	02	0.6	1.5	40	B
19	<i>Terminalia bellirica</i> (Gaertn.) Roxb.	-	3	-	-	2	05	02	1.0	2.5	40	B
20	<i>Terminalia cuneata</i> Roth	-	-	-	1	1	02	02	0.4	1.0	40	B

On the basis of percentage values various species are distributed into five classes, according to Raunkiaer (1934).

**Table 2: Raunkiaer’s percentage values various species.**

Sr. No.	Frequency of plants (in %)	Classes
1	0-20	A
2	21-40	B
3	41-60	C
4	61-80	D
5	81-100	E



**Fig 1: Frequency class and Frequency percentage and compare with Raunkiaer’s value.**

X-axis: Percentage of total number of species. Y-axis: Frequency Classes.

**Table 3: Percentage values various species.**

Frequency class	Class Value	Raunkiaer's value	Frequency class Vegetation
A	0-20	53	20%
B	21-40	14	35%
C	41-60	09	25%
D	61-80	08	15%
E	81-100	16	5%

## Conclusion

The Frequency value refer to the values of Raunkiaer's formula  $A>B>C<D<E$ . The current study shows the frequently occurring species and dominating species found in this sacred grove. The present ecological study of medicinal plant species shows that the given vegetation frequency does not match with Raunkiaer's normal frequency hence vegetation found in Dongargan sacred grove is heterogeneous.

### Acknowledgements:

Author Thete AM thankful to the Principal of the Shri. Dhokeshwar College and Sharma PP is thankful to the management of the College for constant support and encouragements.

**Conflicts of interest:** The authors stated that no conflicts of interest.

## References

- Gadgil M, Berkes F and Folke Carl, "Indigenous Knowledge for biodiversity conservation." *Ambio*, 1993; XII (2-3):266-270.
- Cutis JT. and McIntosh RP, The interrelations of certain analytic and synthetic Phytosociological characters. *Ecology*, 1950; 31:434-455.
- Pradhan SG and Singh NP, Flora of Ahmednagar District. (M.S), Bishen Singh Mahendra Pal Singh. Dehradun, 1999; 339.
- Gadgil M. and Vartak VD, The sacred groves of Western Ghats in India. *Economic Botany*, 1976; 30:152-160.
- Gadgil M. and Vartak VD, Sacred groves of Maharashtra: An inventory. In: SK. Jain (ed) *Glimpses of*

*Indian Ethnobotany*, Oxford and IBH Publishers, New Delhi, India, 1980; 289-294.

- Minakshi Mahajan\* and Sumia Fatima\*\*, Frequency, Abundance and Density of plant species by List count Method, *Epitome: International Journal of Multidisciplinary Research*, 2017; III 21-28.