

Production of citric acid by fermented broth method

Survase Damu Mokinda

Department of Botany, Karmveer Ramraoji Aher Arts, Science & Comm. College, Deola, Tal-Deola, Dist-Nashik, Maharashtra- 423102 (India).

Email: damusurvase@gmail.com | Mob.9284190439

Manuscript Details

Available online on <https://www.irjse.in>
ISSN: 2322-0015

Editor: Dr. Arvind Chavhan

Cite this article as:

Survase Damu Mokinda. Production of citric acid by fermented broth method. *Res. Journal of Science & Engineering*, 2020, Special Issue A9: 133-136.

Article published in Special issue of International e-Conference on "Emerging trends and Challenges In life sciences" organized by Department of Botany, Indraraj Arts, Commerce & Science College, Sillod-431112, Dist Aurangabad, Maharashtra, India date, June 18-19, 2020.



Open Access This article is licensed under a Creative Commons Attribution 4.0 International License, which permits use, sharing, adaptation, distribution and reproduction in any medium or format, as long as you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons license, and indicate if changes were made. The images or other third party material in this article are included in the article's Creative Commons license, unless indicated otherwise in a credit line to the material. If material is not included in the article's Creative Commons license and your intended use is not permitted by statutory regulation or exceeds the permitted use, you will need to obtain permission directly from the copyright holder. To view a copy of this license, visit <http://creativecommons.org/licenses/by/4.0/>

Abstract

Citric acid demand increased in the different industries (fermented food items, beverages and pharmaceutical industries). Ancient source of citric acid was citrus fruits which contain 6 to 9% of citric acid. It was obtained by the processing of citrus fruits and precipitation of calcium salt. Traditional method citrus fruits could not sufficient for fulfill the growing demand of citric acid production. Hence alternative source of citric acid is a microbial origin. Many microorganisms are involved in the production of citric acid. In the present study *Aspergillus niger* is considered more suitable than the other microorganisms for production of citric acid by fermented broth method. Many factors are affecting in vitro production of citric acid. It was confirmed from tabular reading that sucrose was a suitable at 13 % to 15 % carbon source. Many factors were affecting fungal origin production of citric acid like strain of microorganism, temperature, carbon source, PH etc. were considered. Pure strain of *Aspergillus niger* was obtained from soil samples from different localities and from lemon fruits, 2% methanol added into culture medium gives higher yield, 13% sucrose was better carbon source than the glucose, 4.5 P^H acidic was suitable for healthy production of citric acid in vitro condition. Healthy growth was found at 28-30⁰ C temperature.

Keywords: Soil, lemon fruit, Citric acid, Fermentation broth, *Aspergillus niger* etc.

Introduction

Citric acid is widely used an acidifying agent and antioxidant in a fermented food items, beverages and pharmaceutical industries [1]. Microbial fungal origin Citric acid was obtained by submerged fermentation of *Aspergillus niger*. The yield of citric acid depends upon composition of culture medium and on the microbial strain [2].

Citric acid production using *A. niger* is influenced by the process variables such as initial sucrose concentration, pH, nutrient concentration, additive, incubation period, temperature etc. However, demand for Citric acid production is increasing faster than its production and hence requires more economical process. The fitness of solution would be measured by determining the total weight of the proposed solution. Healthy growth of fungus means more quantity of citric acid [3].

In the present study screened different factors for the microbial origin of citric acid like methanol, strain of microorganism, carbon source, PH etc. were considered. Isolation and maintained pure cultures of *Aspergillus niger* from soil sample collected from different localities, suitable 2% methanol added into culture medium gives higher yield, 13% sucrose was gives higher yield than the glucose, 4.5 PH was found more suitable for production of citric acid in vitro and in 28-30° temperature healthy growth was found [4].

Methodology

Aspergillus niger is a fungus recommended for the production of various metabolites. Citric acid is one of the important organic acids synthesized and released on synthetic medium by this fungus. Natural source of *Aspergillus niger* is a soil, hence soil from different localities from Nashik District was collected and tested for higher yielding strain of *Aspergillus niger* for the production of citric acid. It also isolated from lemon fruits.

1. Screening of *Aspergillus niger* for citric acid production:

Selection and isolation of micro-organism which produces high amount of citric acid. Primary screening determines which microorganism is able to produce a citric acid followed by secondary screening to determine capacity of that organism producing quantitatively [5].

2. Isolation of Microorganism:

The natural source of *Aspergillus niger* is the soil. Collection of soil samples from different locations with respect to PH of the soil. In this way ten soil samples

were collected and diluted 1/100, 1/1000 and 1/10000 of each sample. The soil samples were diluted, analyzed and they were purified by sub culturing on Czepecks Dox agar slants. Culture Plates were incubated at 30 °C and isolated cultures were observed and they were purified by sub culturing on Czepecks Dox agar slants. Isolation of *A. niger* from lemon fruits and pure culture isolated and maintained. Similar to soil screening lemon strain tested for citric acid production.

Similarly, fresh and healthy lemon fruits were soaked in the water placed in closed container for a 5- 7 days. After incubation pure cultures of *Aspergillus niger* maintained on culture medium.

3. Screening for organic acid production:

Spore from slant cultures were inoculated on sterile Czepeck's Dox agar medium plates incorporated with Bromo-cresol green dye. Inoculated plates were incubated at 28°C for one to two days and checked colour change blue to yellow indicates organic acid production.

To study the effect of media component on citric acid fermentation, all the components except one to be studied are kept constant with respect to the control medium and one component concentration is changed in particular range. Fermented broths were analyzed for the production of citric acid by selected strains of *Aspergillus niger*.

Results and Discussions

Table 1 Soil from various areas were collected and different cultures isolated from it. These were checked for citric acid production. Similarly different cultures of *A. niger* isolated from lemon and college campus garden soils. The soil pH was checked of the collected soils. Samples were tested for citric acid production and results were noted [6]. The highest yield was reported from LF1, FS3 and CC5. It was found that the highest yield was found when the pH of soil was 4.5, above this pH the yield was less. These three higher yielding strains were used for further experiments [7].

Table No.1: Screening for citric acid yield and selection of strains

Sr. No.	Area	PH	Culture	Yield mg/ml
1.	Lemon fruits		LF1	4.7
			LF2	3.7
			LF3	4.3
			LF4	2.8
			LF5	3.9
2.	Forest soil	6.2	FS1	2.5
		6.1	FS2	3.2
		5.9	FS3	4.1
		5.4	FS4	2.9
		6.1	FS5	3.5
3.	College Campus garden Soil	4.8	CC1	2.6
		5.2	CC2	2.7
		4.9	CC3	1.6
		5.1	CC4	1.8
		4.8	CC5	2.9

Table No.2: Effect of different carbon source % on the production of CA by *Aspergillus niger*

Carbon source	Yield mg/ml Sucrose			Yield mg/ml Glucose			Yieldmg/ml Galactose			Yield mg/ml Fructose		
	LF1	FS3	CC5	LF1	FS3	CC5	LF1	FS3	CC5	LF1	FS3	CC5
%												
10	1.6	2.1	1.9	1.4	1.1	1.3	1.1	0.8	0.8	1.2	1.6	1.7
11	1.9	1.8	1.9	1.4	1.6	1.3	0.5	0.8	0.9	1.1	1.5	1.3
12	2.5	2.6	2.8	1.8	1.6	1.9	0.9	0.7	1.0	1.2	1.5	1.4
13	4.7	3.9	3.6	2.9	2.4	1.8	0.6	0.9	1.6	1.6	1.8	1.2
14	3.1	2.9	2.6	2.4	2.1	2.3	1.1	1.5	1.7	1.9	2.0	1.6
15	2.9	2.4	2.1	1.9	1.8	1.8	0.9	0.8	0.7	1.7	1.9	1.4
16	1.8	1.4	1.3	1.0	1.2	1.3	0.5	0.7	0.6	0.9	0.9	0.8

Table No. 3: Effect of pH on the production of CA by *Aspergillus niger*

Sr. No.	pH	LF1	FS3	CC5
1.	2.5	2.6	2.8	2.9
2.	3.0	2.8	2.8	3.0
3.	3.5	3.0	3.1	3.3
4.	4.0	2.9	3.5	3.9
5.	4.5	3.9	3.7	3.8
6.	5.0	1.8	2.1	1.9
7.	5.5	1.3	1.2	1.5

Table No.4: Ascending chromatography of organic acid and test solution

Sr. No.	Sugar	Distance travelled by solvent	Distance travelled by solute	Rf	% Rf
1.	Citric acid	10.0	4.0	0.4	40
2.	Oxalic acid	10.0	5.0	0.5	50
3.	Succinic acid	10.0	6.0	0.6	60
4.	Test	10.0	4.0	0.4	40

Table2, when all the three cultures were tested for variation in sugar concentration (Sucrose and glucose). It was found that glucose is not a suitable carbon source for citric acid production. Culture medium with sucrose showed higher yields. It was reported that at a sugar concentration of 13% all the three cultures gave the higher yield. It was also found that at a sugar concentration of (15% to 18%) the yield reported was less as compared to 13%. Kovats [8] reported that higher sugar concentration (15 to 18%) greater amount of residual sugars remains in the medium and process become uneconomical.

Table3, pH plays an important role in the production of citric acid *in vitro*. A pH of (2.5 to 5.5) was tested for citric acid production. It was found that a pH of 4.5 was found to be suitable for all the three cultures for the yield of citric acid [9]. This showed a contrast with the finding of Prescott and Dunn [10] which claims that the initial pH for sucrose.

Table4, Ascending chromatography was done by using the solvent system n-Butanol, formic acid and water in the proportion of 10:2:5 respectively. It was prepared by separating funnel and out of the two layers upper organic layer was used. The fermented broth concentrated by evaporating it in a Petri dish. Chromatogram removed, dried and sprayed with 0.4% Bromo-Cresol Green prepared in ethanol (pH is equal to 6.7) and R_f values were calculated and recorded. Calculated R_f value of samples were compared with the different organic acids and from this it was confirmed that fermentation broth contains citric acid

Conflicts of interest: The authors stated that no conflicts of interest.

References

- Currie JN. The citric acid fermentation of *Aspergillus niger*, *Can. Journal of microbiology*, 1917,7, 447-453.
- Aftab Nadeem, Quratulain Sayed, Shahjahan Baig, Muhammed Irfan and Muhammed Nadeem. Enhanced production of citric acid by *Aspergillus niger* M-101 using lower alcohols, *Turk Biyokimya Dergisi, Turkish Journal of Biochemistry*, 2010, 35; 10; 7-13
- K. Anand Kishor, M. Praveen Kumar, V. Ravi Krishna and G. Venkat Reddy(2008): Optimization of process variables of citric acid production using *Aspergillus niger* in a batch fermenter, *Engineering Letter*, 16:4, EL_4_17.
- Chaturvedi Madhusudan, Singh Manoj, Chugh M Rishi (2010): Citric acid production from cane molasses using submerged fermentation by *Aspergillus niger*. *ATCC9142, Journal of Pharmacy Research* 2010, 3(6), 1215-1222.
- Abdullah - Al -Mahin, Shek Mehdi Hasan ,MahboobHossain Khan and Rehan Begum (2004) : Citric Acid production by *Aspergillus niger* through solid state cane bagasse, *Bangladesh ,Vol 25 ,Number 1, June 2008 , pp-9-12.*
- SING, S.P.; VERMA, U.N.; KISHOR, M. and SAMDANI, H.K. (1998): Effect of medium concentration on citric acid production by submerged fermentation. *Orient Journal of Chemistry*, March 1998, vol. 14, no. 1, p. 133-135
- Ashish Kumar and V.K. Jain (2008): Solid state fermentation studies of citric acid production, *African Journal of Biotech* Vol. 7 (5) pp. 644-650.
- Kovats (1960): Studies on submerged citric acid fermentation, *Acta microbiology Pat* 9, 275-287.
- Gupta J.K., Hedging L.G. & Jorgensen O.B. (1976): Effect of sugars, pH and Ammonium nitrate on formation of Citric acid by *Aspergillus niger*, *Acta. Microbiology Acad. Science Hung.* 23, 63-67.
- Prescott and Dunn (1987): *A Industrial microbiology*, 4th edition, CBS publisher and distributor, New Delhi, India, August, 1987, p 710-715.