

Effect of *Boswellia serrata* gum on seed germination of some seeds.

Badar KV

Department of Botany, Yeshwantrao Chavan College, Sillod, Dist. Aurangabad (M. S.) India

Email : drkvbadar@gmail.com

Manuscript Details

Available online on <https://www.irjse.in>
ISSN: 2322-0015

Editor: Dr. Arvind Chavhan

Cite this article as:

Badar KV. Effect of *Boswellia serrata* gum on seed germination of some seeds., *Int. Res. Journal of Science & Engineering*, 2020, Special Issue A9: 103-106.

Article published in Special issue of International e-Conference on "Emerging trends and Challenges In life sciences" organized by Department of Botany, Indraraj Arts, Commerce & Science College, Sillod-431112, Dist Aurangabad, Maharashtra, India date, June 18-19, 2020.



Open Access This article is licensed under a Creative Commons Attribution 4.0 International License, which permits use, sharing, adaptation, distribution and reproduction in any medium or format, as long as you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons license, and indicate if changes were made. The images or other third party material in this article are included in the article's Creative Commons license, unless indicated otherwise in a credit line to the material. If material is not included in the article's Creative Commons license and your intended use is not permitted by statutory regulation or exceeds the permitted use, you will need to obtain permission directly from the copyright holder. To view a copy of this license, visit <http://creativecommons.org/licenses/by/4.0/>

Abstract

The gum of *Boswellia* species has been used as incense in religious and cultural ceremonies and in medicines since time immemorial. Gum is naturally occurring chemical substance in the plant, *Boswellia serrata* (*Salai/Salai guggul*), is a moderate to large sized branching tree of family Burseraceae (Genus *Boswellia*), grows in dry mountainous regions of India, Northern Africa and Middle East. After processing, the gum is then graded according to its flavour, colour, shape and size. In India, the States of Andhra Pradesh, Gujarat, Maharashtra, Madhya Pradesh, Jharkhand and Chhattisgarh are the main source of *Boswellia serrata*. There are different name of plants according to locality. Gum is a substance which exudates naturally from the stem or from the wounds of trees. Gums are colloidal in nature soluble in water but completely insoluble in alcohol and ether. Impact of gum collected from was studied for different time period of 1% gum concentration on the germination of cereals, pulses and oil seeds like wheat, jowar, cowpea, gram, safflower and soybean. Percent germination or percent inhibition of germination, root and shoot length of seedlings was measured after 7 days of incubation at room temperature. It was concluded from the present investigation that 1% concentration of *Boswellia serrata* gum was found to promote the germination of wheat, jowar, cowpea, and soybean at 10-hour treatment.

Keywords: Gum, *Boswellia*, Seed Germination.

Introduction

Gums are a group of plant product resembling carbohydrates and widely distributed in vegetable kingdom. The vegetable gums constitute an important group of economic plant product and are utilized in several ways. It is very commonly used by human being from ancient time. Now a day in modern industry large quantity of gum have been used for manufacture of pharmaceutical products. Gums are characterized by ability to dissolve in water forming viscid solution by absorbing water to form jelly gelatinous paste. When these jelly exposed to air these gelatinous paste loose the water and dry to hard clear or with different color Gums are characterized by ability to dissolve in water forming viscid solution by absorbing water to form gelatinous paste. In some cases the production of gum has been attributed to fungi attacking the plant, these fungi being responsible for enzymes that penetrate the tissues and transform the celluloses and hemicelluloses of the cell wall into gum.

Malcolm [1] concluded that the production of gum in Sudan gum Arabic trees is due to bacterial agency. The real cause of production of gum in many trees is uncertain. The best use of gum is to prepare sticky substance for pasting the paper and other things. Treatment of gums to seeds has positive effect on seed germination. *Acacia* gum is being widely used as an experimental vehicle for drugs in physiological and pharmacological experiments, and it is supposed to be an inert substance, recent reports have confirmed that it has some biological properties as an antioxidant [2-4] on the metabolism of lipids [5-6], positive contribution in treating kidney, [7-9], cardiovascular [10] and gastrointestinal diseases [11-12]. *Acacia* gum is usually referred as Gum Arabic (GA). It is an edible biopolymer obtained as exudates of mature trees of *Acacia* spp. The exudate is a non-viscous liquid, rich in soluble fibers, and its emanation from the stems and branches usually occurs under stress conditions such as drought, poor soil fertility, and injury [13].

Gums are to be found in a greater or less degree in most plant families. The gum occurs in about 44 families covering 1,900 genera and 21,000 species. Various organs of the plant may produce or secrete them. They may be produced only in very small quantity and not be readily discernible or they may be produced very copiously forming large, conspicuous incrustations on the surface, as with most of the commercial gums, particularly the tree gums. Certain families of flowering plants are notable for the number of species they contain that are free gum yielders. Among them notable is the Leguminosae, in which a hundred or more species of *Acacia* alone are known to yield gum, including those that are commercially important for gum arabic. Several species of *Astragalus* are also free gum yielders and are the source of gum tragacanth. Some additional notable gum yielding genera in the family are *Albizia*, *Bauhinia*, *Caesalpinia*, *Ceratonia* and *Pithecolobium*. Other important gum yielding families are Anacardiaceae, Combretaceae, Meliaceae, Rosaceae and Rutaceae.

Methodology

Collection of Gum samples:

Plant gums were regularly collected in all the seasons. It was done by using axe, sterilized blade. Fine cut was made at different parts of the plant, like root, stem, leaves, flower and fruits. Later on at 30, 45 and 60 days exudates gums were collected in presterilized plastic bags, kept in laboratory condition until it was used [14].

Preparation of fine powder of Gum:

The fine powder of collected dry gums was prepared by using mixer grinder and kept in clean glass pots at 4-6°C temperature.

Application of gum on seed germination:

The effect of different time period (1 hr, 3 hr, 5hr, 10hr) of plant gum on seed germination and the impact of different concentration like 1%, 2%, 3% and 4% on cereal, pulses and oil seeds of seed mycoflora have been studied. The impact of different time period of percent gum concentration of *Boswellia serrata* were studied on the germination of cereals,

pulses and oil seeds like wheat, jowar, cowpea, gram, safflower, soybean and the results are summarized in Table. *Boswellia serrata* gum was applied on germinating seeds for 1 to 10 hour treatments. The results were recorded at different time intervals by measuring shoot length and root length by vigour formula [15].

Results and Discussions

It was observed from the 1% concentration of *Boswellia serrata* gum was found to promote the germination of wheat, jowar, cowpea and soybean at 10 hour treatment. 1 hour treatment was found to promote the germination of wheat, cowpea, gram and soybean.

Table 01: Impact of different time period of *Boswellia serrata* gum on seed germination (1% gum concentration)

Sr. No.	Seed plant Name	1 hr			3 hrs			5 hrs			10 hrs			Control		
		% germ	Shoot length (mm)	Root length (mm)	% germ	Shoot length (mm)	Root length (mm)	% germ	Shoot length (mm)	Root length (mm)	% germ	Shoot length (mm)	Root length (mm)	% germ	Shoot length (mm)	Root length (mm)
1.	Wheat	80	4.2	2.2	70	4.0	3.1	80	4.9	2.7	90	6.0	5.2	70	3.4	2.1
2.	Jowar	70	7.2	5.5	80	3.9	3.0	70	5.3	2.8	90	5.7	4.2	80	6.1	5.9
3.	Cowpea	80	3.2	3.2	70	3.7	2.9	90	4.7	3.8	70	4.3	3.8	60	2.9	2.8
4.	Gram	80	3.2	2.3	80	4.2	2.7	80	5.0	3.2	60	4.1	3.4	60	1.8	2.3
5.	Safflower	70	3.5	2.7	80	4.1	3.3	80	5.2	3.1	70	3.9	3.6	70	3.1	2.6
6.	Soyabean	70	3.6	3.5	80	3.8	2.9	80	5.1	2.9	80	3.1	3.8	80	3.4	2.9

It was also found to promote the shoot length, root length of wheat, cowpea, gram whereas it was found to retard the growth of shoot and root length of jowar and soybean. Similar type of results was recorded for 3-hour 5-hour treatment. Whereas, at 10-hour treatment the shoot and root length of all treated seeds except Soyabean were found to be increased as compared to control.

The *Boswellia serrata* gum is being widely used for industrial purposes such as a stabilizer, a thickener, an emulsifier and an encapsulating in the food industry, and to a lesser extent in textiles, ceramics, lithography, cosmetic and potential Anti-inflammatory agent [16] Hence, the *Boswellia serrata* gum can be useful for induces maximum seed germination in various ways.

Conflicts of interest: The authors stated that no conflicts of interest.

References

1. Malcolm DW. Report on Gum and Gum arabic (Dar-es-salaam Gov. Printer). 1936, 3(4): 188-191.
2. Trommer H and Neubert RH. The examination of polysaccharides as potential antioxidative compounds for topical administration using a lipid model system. *International Journal of Pharmaceutics*, 2005, 298(1):153-163.
3. Ali, B.H. and M.S. Al Moundhri. Agents ameliorating or augmenting the nephrotoxicity of cisplatin and other platinum compounds: a review of some recent research. *Food and Chemical Toxicology*, 2006, 44(8): 1173-1183.
4. Hinson, J.A., A.B. Reid, S.S. McCullough and L.P. James. Acetaminophen-induced hepatotoxicity: role of metabolic activation, reactive oxygen/nitrogen species, and mitochondrial permeability transition. *Drug Metabolism Reviews*, 2004, 36(3-4): 805-822.
5. Tiss, A., F. Carrière and R. Verger. Effects of gum arabic on lipase interfacial binding and activity. *Analytical Biochemistry*, 2001, 294(1): 36-43.

6. Evans, A.J., R.L. Hood, D.G. Oakenfull and G.S. Sidhu. Relationship between structure and function of dietary fibre: a comparative study of the effects of three galactomannans on cholesterol metabolism in the rat. *British Journal of Nutrition*, 1992, 68(1): 217-229.
7. Matsumoto, N., S. Riley, D. Fraser, S. Al-Assaf, E. Ishimura, T. Wolever, G.O. Phillips and A.O. Phillips. Butyrate modulates TGF-beta1 generation and function: potential renal benefit for *Acacia* (sen) SUPERGUM (G.A.)? *Kidney International*, 2006, 69(2): 257-265.
8. Bliss, D.Z., T.P. Stein, C.R. Schleifer and R.G. Settle. Supplementation with GA fiber increases fecal nitrogen excretion and lowers serum urea nitrogen concentration in chronic renal failure patients consuming a low-protein diet. *The American Journal of Clinical Nutrition*, 1996, 63(3): 392-398.
9. Ali, A.A., K.E. Ali, A. Fadlalla and K.E. Khalid. The effects of GA oral treatment on the metabolic profile of chronic renal failure patients under regular haemodialysis in Central Sudan. *Natural Product Research*, 2008, 22(1): 12-21.
10. Glover, D.A., K. Ushida, A.O. Phillips and S.G. Riley. *Acacia*(sen) SUPERGUM™ (Gum arabic): An evaluation of potential health benefits in human subjects. *Food Hydrocolloids*, 2009, 23(8): 2410-2415.
11. Wapnir, R.A., B. Sherry, C.N. Codipilly, L.O. Goodwin and I. Vancurova. Modulation of rat intestinal nuclear factor NF-kappaB by gum arabic. Rat small intestine by gum arabic. *Digestive Diseases and Sciences*, 2008, 53(1): 80-87.
12. Rehman, K.U., M.A. Wingertzahn, S. Teichberg, R.G. Harper and R.A. Wapnir. Gum arabic. (GA) modifies paracellular water and electrolyte transport in the small intestine. *Digestive Diseases and Sciences*, 2003, 48(4): 755-760.
13. Williams, P.A., G.O. Phillips and A.M. Stephen. Spectroscopic and molecular comparisons of three fractions from *Acacia senegal* gum. *Food Hydrocolloids*, 1990, 4(4): 305-311.
14. Badar, K.V. Studies on antimicrobial properties of gum of some plants. Ph.D. Thesis, Dr. Babasaheb Ambedkar Marathwada University, Aurangabad, 2011.
15. Abdul- Baki A.A, and Anderson J.D. Vigour dermination in soyabean seed by *multiple criteria crop sci. B* 1975, 630-633
16. Siddique M.Z. *Boswellia serrata*, a potential Anti-inflammatory agent: An Overview. *Indian Jouurnal of Pharmaceutical sciences*, 2011, 73 (3) :255-261.