REVIEW ARTICLE OPEN ACCESS

Phytochemical studies on stem bark of Ceiba pentandra linn

Revansiddha Dhotre and Mamtaram Kare

Department of Botany, Pratishthan Mahavidyalaya, Paithan, Tal: Paithan Dist: Aurangabad Pin: 431107. Email: revan945@gmail.com

Manuscript Details

Available online on https://www.irjse.in
ISSN: 2322-0015

Editor: Dr. Arvind Chavhan

Cite this article as:

Revansiddha Dhotre and Kare Mamtaram. Phytochemical studies on stem bark of *Ceiba pentandra* linn., *Int. Res. Journal of Science & Engineering*, 2020, Special Issue A9:236-240.

Article published in Special issue of International e-Conference on "Emerging trends and Challenges In life sciences" organized by Department of Botany, Indraraj Arts, Commerce & Science College, Sillod-431112, Dist Aurangabad, Maharashtra, India date, June 18-19, 2020.

Open Access This article is licensed under a Creative Commons Attribution International License, which permits use, sharing, adaptation, distribution and reproduction in any medium or format, as long as you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons license, and indicate if changes were made. The images or other third party material in this article are included in the article's Creative Commons license, unless indicated otherwise in a credit line to the material. If material is not included in the article's Creative Commons license and your intended use is not permitted by statutory regulation or exceeds the permitted use, you will need to obtain permission directly from the copyright holder. To view a copy of this license, visit http://creativecommons.org/ licenses/by/4.0/

Abstract

Ceiba pentandra. is one of such ayurvedic remedy that has been mentioned in many Indian medicinal literatures. It is crucially significant as it has the largest number of phytochemicals and secondary metabolites pharmacological and biological properties. The present work deals development and standardization phytochemical analysis for quantification of stem bark extract of medicinal plant of Ceiba pentandra Linn. The scientific parameter is necessary to identify the exact plant material and to find its quality and purity. The present study deals with various preliminary phytochemical analyses of various successive extracts such as qualitative chemical analysis. Tannins, saponins, alkaloids and flavonoids content of stem bark of the title plant have been determined. These studies indicated the possible information for correct identification and standardization of this plant material.

Keywords: *Ceiba pentandra,* Phytochemical Analysis, Bark Extract, Solvents.

Introduction

There are a number of floras in use for medicinal purposes over the past several centuries. Countries such as China, India, and Egypt are well known for the active usage of medicinal plants in the treatment of various incurable diseases. India is the largest producer of medicinal herbs in the world due to which it is often called a botanical paradise. Ayurvedic science is deeply rooted in India and its neighboring countries. It was developed even before the medieval period, when people had little knowledge of science. There are a number of ancient therapeutic measures based on medicinal plants that have been developed in India

Dhotre and Kare, 2020 237

[1-4] . They can cure several diseases and ailments such as diabetes, cardiovascular disorders, cancer, and liver damage [1,5,6,7].

Tree bark is very complex in structure and has the potential of containing many primary and secondary metabolites. Products stored in the bark are useful for preparation of many drugs. The complex structure of the bark can be utilized for botanical identification to maintain the quality and purity of the drug [8]. In the present study, attempts will be made on barks of medicinal trees which are available in Marathwada region by applying various techniques and parameters such as Morphology, Anatomy, Maceration, Pharmacognosy and Phytochemistry. The aim of present work is to develop the parameters to detect the genuinity of bark drugs, as the authentic material is prerequisites for the preparations of home remedies and for industrial use parameters to detect the genuinity of bark drugs, as the authentic material is prerequisites for the preparations of home remedies and for industrial use.

Ceiba pentandra recently the plant has undergone extensive scientific research for its medicinal value. In Ayurvedic its bark decoction has been used as a diuretic, aphrodisiac, and to treat headache, as well as type II diabetes, in many countries in Africa, the bark is taken as remedies for diarrhoea, localized oedemas, wash sores, furuncles leprous macules, to relieve stomach complaints, hernia, blennorrhoea, heart-trouble, asthma, gargles for gingivitis, aphtes and sometimes toothache, also used in India and Malaya for bowel-complaint and West Africa for diarrhea, In Nigeria the bark infusion is used for febrifuge [9]. Nigeria it is used for bath, barkdecoction are considered excellent for curing evening fever especially those deemed to arise from evil influence [10]. The stem bark is used locally as myriad of effects on medical conditions such as treatment of wounds, cough, high blood pressures, diarrhoea, dysentery, yellow fever and tumours [11].

Tail trees with smooth, green stem and horizontally spreading whorled branches. Leaves digitate; leaflets 5-9, lanceolate, 5-12 cm long, cuspidate, green above, glaucous beneath; petioles 10-15 cm long; petiolules 2-3

mm long. Flowers 3-5 cm across, clustered in the leaf axils; pedicels 2-4 cm long. Calyx 1.2-2 cm long,5-lobed, glabrous outside, hairy. within; lobes triangular-ovate. Corolla dull white or pinkish; petals obovate-oblong, 3.5-4 cm long, woolly outside. Anthers sinuous. Ovary conical, 1.5-2 cm long, glabrous. Capsules fusiform, 3.5-4 cm long, hairy. Seeds pyriform, 5-6 mm long, black, embedded in silky wool.

Materials and Methods

Plant Material

The stem bark of *Ceiba pentandra* was collected from Pratishthan Mahavidyalaya, Paithan. The plant materials was identified and Authenticated by Dr. M. A. Kare, Department of Botany, Pratishthan Mahavidyalaya, Paithan.

Preparation of Extract

The stem bark of Ceiba pentandra shaded dried, and then these are made into coarsely powdered form using dry grinder. The powdered bark of the plant (180gm.) was packed in soxhlet apparatus and continuously extracted with petroleum ether (40-600C) till complete extraction, after completion of extraction the solvent was removed by distillation and then concentrated extract obtained was dried under reduced pressure using rotatory evaporator at temperature not exceeding 400C and then give moderate heating on water bath. A yellowish extract approximate 1 gm. was obtained. From the drug petroleum ether was removed and the defatted drug was extracted with methanol (95%) till complete extraction, after completion of extraction the solvent was removed by distillation and then concentrated extract obtained dried under reduced pressure at temperature not exceeding 400C and then give moderate heating on water bath. The methanolic extract obtained was dark yellow in color, weighed about 42.8 gm. The methanolic extract was kept in Petridis and it was stored in desiccators at cool place [12].

Results

Phytochemical test of Ceiba pentandra

Table 1: Observation of Quantitative analysis of organic of Ceiba pentandra

Sr. No.	Quantitative Standards	%w/w
1	Total ash	7.12
2	Acid soluble ash	5.80
3	Acid insoluble ash	1.32
4	Water soluble ash	4.28
5	Water insoluble ash	1.34
6	Loss of weight on drying 105°C	52.8
7	Alcohol soluble extractive value	6.73
8	Water soluble extractive value	6.32

Table 2: Physico-Chemical Properties of Ceiba pentandra . stem bark.

Test	Methanolic Extract	Petro. Ether	Chloroform	Acetone
Tannins	+	-	+	+
Phenols	+	-	-	-
Alkaloids	+	-	-	-
Saponins	+	-	-	-
Iridoids	-	+	-	+
Quercetin	-	-	-	-
Kaempferol	-	-	-	-
Catechin	-	-	-	-
Coumarin	-	+	+	+
6,7-Dimethoxy	-	-	+	+
coumarin				
5-Methoxy genistein	-	-	+	-
Anthocyanin	-	+	-	-
Proanthocyanin	-	+	-	-
Carbohydrates	+	-	-	+
Flavonoids	+	-	+	-
Glycosides	+	-	+	-
Proteins	-	-	+	+

Table 3 Successive Extractive Values of the stem Bark of Ceiba pentandra.

Sr. No.	Solvent	Weight of Drug	Average Extractive Value (%w/w)
1	Methanol	10gm	5.4
2	Alcohol	10gm	13.1
3	Benzene	10gm	6.8
4	Petroleum	10gm	1.09
5	Chloroform	10gm	2.90
6	Acetone	10gm	4.21

Dhotre and Kare, 2020 239

In plants phytochemicals are naturally present. They give colour, flavor, smell and texture. A part from that, phytochemicals could prevent diseases including cancer and cardiovascular diseases and inhibit pathogenic microorganisms. Nowadays the use of medicinal plants rapidly increases in medicine [12]. Phytochemical evaluation of methanolic extract of Ceiba pentandra showed the presence of Tannins, Phenols, Alkaloids, Saponins, Iridoids, Quercetin, Kaempferol, Catechin, 6,7-Dimethoxy Coumarin, coumarin, 5-Methoxy genistein, Anthocyanin, Proanthocyanin, Carbohydrates, Flavonoids, Glycosides and Proteins.

Phytochemical screening of the bark showed some differences in the presence of phytoconstituents which are known to have importance in medicine [13,14,15,16,17,18] The preliminary screening tests may be useful in the detection of the bioactive principles and subsequently may lead to the drug discovery and development [19,20, 21,22]

Saponins are found only in petroleum ether extract. Chloroform extract revealed the presence carbohydrates and saponins. Ethanol extract showed the presence of alkaloids, flavonoids, glycosides, phenols, steroids, terpenoids and quinones whereas acetone extract showed the availability of alkaloids, flavonoids, phytosterols, phenols and proteins. Such alkaloids were effective against ovarian, brain, breast, lung cancer etc. [23,24,25,26] and several of its semisynthetic analogues are 9-Nitro-CPT, 10-hydroxy-9-dimethylaminomethyl -CPT, 7-Ethyl- 10 -hydroxy-camptothecin (SN-38), are applied as clinical anticancer drugs in USA, Europe and Japan [27]. Other alkaloids include indicine, indicine Noxide, thalicarpine and tetrandrine [28]. Flavonoids are also reported to have inhibitory action on growth and proliferation of different types of tumors [29].

Conclusion

Various extracts was subjected to Pharmcoganostic Evaluation for the identification of various Phytoconstituents and rest of extracts were utilized for pharmacological screening. Phytochemical evaluation extract of Ceiba pentandra showed the presence of Tannins, Phenols, Alkaloids, Saponins, Catechin, Quercetin, Kaempferol, Coumarin, 6,7coumarin, 5-Methoxy Dimethoxy genistein, Anthocyanin, Proanthocyanin, Carbohydrates, Flavonoids, Glycosides and Proteins.

Acknowledgement:

The authors are grateful to Principal, Pratishthan Mahavidyalaya, Paithan, for providing the necessary laboratory facilities and we are also thankful with our deepest core of heart to Dr. M. A. Kare for his valuable guidance.

Conflicts of interest: The authors stated that no conflicts of interest.

References

- 1. Dwevedi, Alka, Sharma, Kuldeep, Yogesh, K, Sharma. "Cadamba: A miraculous tree having enormous pharmacological implications." *Pharmacognosy Reviews* 9, no. 18 (2015): 107-113.
- Zaidan MR, Noor Rain A, Badrul AR, Adlin A, Norazah A,. "In vitro screening of five local medicinal plants for antibacterial activity using disc diffusion method." *Trop Biomed* 22 (2005): 165-170.
- 3. Ahmad I, Mehmood Z, Mohammad F. "Screening of some Indian medicinal plants for their antimicrobial properties." *J Ethnopharmacol* 62 (1998): 183-193.
- 4. Bhakuni DS, Dhar ML, Dhar MM, Dhawan BN, Mehrotra BN. "Screening of Indian plants for biological activity." *Indian J Exp Biol* 7 (1969): 250-262.
- 5. Ríos JL, Recio MC. "Medicinal plants and antimicrobial activity." *J Ethnopharmacol* 100 (2005): 80-84.
- 6. Alekhya V, Deepan T, Sahoo S, Dhanaraju MD. "Preliminary phytochemical screening and evaluation of in vitro antioxidant activity of Anthocephalous cadamba by using solvent extracts." *Eur J Biol Sci* 5 (2013): 34-37.
- 7. Kirana C, McIntosh GH, Record IR, Jones GP. "Antitumor activity of extract of Zingiber aromaticum and its bioactive sesquiterpenoid zerumbone." *Nutr Cancer* 45 (2003): 218-225.
- Usman, S.B., Kyari, S.U., Abdulrahman, F.I., Ogbe, A.O., Ahmad, G.Y., Ibrahim, U.I., Sakuma, A.M. (2012): Proximate composition, phytochemical and elemental analysis of some

- organic solvent extract of the wild mushroom-Ganoderma lucidum. J. Nat. Sci. Res., 2:24-35.
- Burkill, H.R.M. The Useful Plants of West Tropical Africa, vol. 1 (London: Royal Botanic Gardens, Kew), 1985.
- 10. Hutchinson, J. and Dalziel, J.M. (1954) Flora of West Tropical Africa. Vol. I, Part 1, 2nd Edition, Agents for Overseas Governments Administrations, London.
- 11. Sandhya, S., Sai Kumar, P., K.R, V., David, B., & Kumar, K. (2011). Plants as Potent Antidiabetic and Wound healing agents. Hygeia Journal of Drug and Medicine, 3(1), 11-19.
- 12. Renu, S. "Useful Metabolites From Plant Tissue Cultures." Biotechnology 4, no. 2 (2005): 79-93.
- 13. Sukumaran S, Kiruba S, Mahesh M, Nisha SR, Miller Paul Z, Ben CP, Jeeva S. "Phytochemical constituents and antibacterial efficacy of the flowers of Peltophorum pterocarpum(DC,) Baker ex Heyne." Asian Pacific Journal of Tropical Medicine 4, no. 9 (2011): 735-738.
- 14. Kiruba S, Mahesh M, Nisha SR, Miller Paul Z, Jeeva S. "Phytochemical analysis of the flower extracts of Rhododendron arboreum Sm.ssp. nilagiricum (Zenker) Tagg ." Asian Pacific Journal of Tropical Biomedicine, 2011: S284-S286.
- 15. Jeeva S, Johnson M, Aparna JS, Irudayaraj V. "Preliminary phytochemical and antibacterial studies on flowers of selected medicinal plants." International Journal of Medicinal and Aromatic Plants 1, no. 2 (2011): 107-114.
- 16. Jeeva S, Johnson M. "Anti-bacterial and phytochemical studies on methanolic extracts of Begonia flaccifera Bedd. flower." Asian Pacific Journal of Tropical Biomedicine 1, no. S1 (2012): S151-S154.
- 17. Johnson M, Aparna JS, Jeeva S, Sukumaran S, Anantham B. "Preliminary phytochemical studies on the methanolic flower extracts of some selected medicinal plants from India." Asian Pacific Journal of Tropical Biomedicine 1, no. S1 (2012): S79-S82.
- 18. AR, Florence, Joselin J, Sukumaran S, and Jeeva S. "Screening of Phytochemical Constituents from Certain Flower Extracts." International Journal of Pharmacy Review & Research 4, no. 3 (2014): 152-159.
- 19. Joselin J, Jenitha S, Brintha TSS, Jeeva S, Sukumaran S, Geetha VS. "Phytochemical and FT-IR Spectral analysis of certain Bamboo species of South India." Biodiversity, Biopropecting and Development 1, no. 103 (2014): 1-2.
- 20. Joselin J, Shynin Brintha TS, Florence AR and Jeeva S. "Screening of select ornamental flowers of the family Apocyanaceae for phytochemical constitutions." Asian Pacific Journal of Tropical Disease 2 (2012): 1-6.
- 21. Florence AR, Joselin J, Jeeva S. "Intra-specific variation of bioactive principles in select members of the genus

- Clerodendrum L." Journal of Chemical and Pharmaceutical Research 4, no. 11 (2012): 4908-4914.
- Joselin J, Shynin Brintha TS, Florence AR, Jeeva S. "Phytochemical evaluation of Bignoniaceae flowers." Journal of Chemical and Pharmaceutical Research 5, no. 4 (2013): 106-111.
- 23. Sato N, Mizumoto K, Kusumoto M, Niiyama H, Maehara N, Ogawa T, Tanaka M. "9-Hydroxyellipticine inhibits telomerase activity in human pancreatic cancer cells." FEBS Lett 44 (1998): 318-321.
- 24. Hsiang YH, Hertzberg R, Hecht S, Liu LF. "Camptothecin induces protein linked DNA breaks via mammalian DNA topoisomerase I." J. Biol. Chem 260 (1985): 14837-14878.
- Chen AY, Liu LF. "Design of topoisomerase inhibitors to overcome MDRI- mediated drug resistance." Advanced Pharmacology, (New York) 29 (1994): 245-256.
- Multon E, Riou JF, Lefevre D, Ahomadegbe JC, Tiou G. "Topoisomerase II- mediated DNA cleavage activity induced by ellipticines on the human tumor cell line N 417." Biochem Pharmacol 38 (1989): 2077-2086.
- 27. Godwin S, Smith AF, Horning EC. "Alkaloids of Ochrosia elliptica." J. Amer. Chem Soc 81 (1959): 1903-1908.
- Hartwell JL, Abbott BJ. "Antineoplastic principles in plants: Recent developments in the field." Adv Pharmacol Chemother 7 (1969): 117-209.
- Netto, CC. "Cranberry and its phytochemicals: A review of In vitro anticancer studies." J. Nutr 137 (2007): 186S-193S.

© 2020 | Published by IRJSE